

A non-hormonal and non-steroidal gel for the management of vulvovaginal symptoms

Ideal for short- and long-term use









Vulvovaginal conditions

Quality of life of many women can be significantly impacted by **vulvovaginal symptoms** such as:

- · Vaginal dryness and burning sensation
- Itching and tenderness
- Painful intercourse (dyspareunia)
- Painful urination (dysuria)
- Bowel movement (defecation)

Most women are often **reluctant or embarrassed to discuss their symptoms** or they assume it is an **expected and natural part of the aging process.**¹

Approximately **70%** of women with the above symptoms **do not discuss** the issue with their Health Care Providers and 40% expect doctors to initiate this conversation.^{1,2}

Vaginal atrophy (VA)

Vaginal dryness is the most prevalent and bothersome symptom as it affects up to 93% of women with VA.³

Affects
up to **90***postmenopausal
women⁴

15% female's experiences symptoms before menopause⁵

up to 70% patients affected by breast cancer present menopausal symptoms²

Laser and radiofrequency devices are used to reduce VA symptoms, and are often referred to as **vaginal rejuvenation**. The most common side effects are burning sensation, discomfort, swelling and redness.⁶

Lichenoid dermatoses

Lichen sclerosus (LS), Lichen planus (LP) and Lichen simplex chronicus (LSC) are chronic and progressive inflammatory diseases characterized by genital skin thinning. They are most prevalent in females **during prepubertal and post-menopausal period.**^{7,8}

Achieving symptom relief and finding an effective treatment for long-term use, without side effects is crucial for both physicians and patients.^{3,5}



StrataMGT® - the innovative gel for the management of vulvovaginal conditions

StrataMGT is a non-hormonal and non-steroidal gel.

StrataMGT **supports** vulvovaginal mucosal conditions and **speeds up** recovery post vaginal rejuvenation.

StrataMGT is a **suitable alternative** to vaginally administered estrogen and topical coricosteroids.⁹

StrataMGT **relieves symptoms** such as itchiness, tenderness, dryness, burning sensation, painful intercourse, painful urination and rectal or defecating pain.¹⁰

StrataMGT **improves** erythema, mucosal tissue thinning, erosions, fissures, ulcerations, scarring/adhesions and swelling.⁹


StrataMGT is indicated for **short- and long-term use without the side effects** of hormonal therapy and topical corticosteroids.

Strata Mgt

Clinical evidence

A clinical study was conducted in Laguna Hills, CA, USA to evaluate the efficacy and safety of StrataMGT and the improvement of patient's Quality-of-Life (QoL).¹⁰

Patients rated the impact of vulvar conditions on QoL using Vulvar Quality of Life Index (VQLI) ranging from 0=not impacted, 1-15=mildly impacted, 16-30=moderately impacted, 31-45=severely impacted.





StrataMGT is **highly efficient in reducing clinical symptoms** associated with VA and improving QoL.


Clinical evidence


A clinical study was conducted in Brisbane, Australia to evaluate the efficacy and safety of StrataMGT for the management of genital skin conditions.⁹

Patients rated severity of the symptoms using a 10-point scale ranging from 0=normal to 10=worst possible.









All symptoms significantly decreased during the first 6 weeks, **including up to a 100% resolution** for some patients.

Continuous **improvements** were seen and maintained after **2 years post start of the treatment.**

StrataMGT is a **suitable alternative treatment** for women suffering from various highly symptomatic vulvovaginal conditions **without the side effects** of topical corticosteroids and vaginally administered hormonal therapy.

StrataMGT® indications

StrataMGT is used to restore the barrier function and moisture balance of the mucosal epidermis.

StrataMGT was developed for use on **all types of mucosal conditions** and post-procedures care including:

- Vaginal atrophy / Atrophic vaginitis
- Genitourinary syndrome of menopause (GSM)
- Vulvovaginitis (including pre-pubertal)
- Vulvodvnia
- Lichenoid dermatoses
- Vaginal grazing and Episiotomy
- Post vaginal rejuvenation
- Incontinence related dermatitis and Diaper rash
- Hemorrhoids and anal fissures
- Allergic and irritant dermatitis



MUCOSAL HEALING

StrataMGT may be directly applied to dry, wet, cracked and sensitive mucosal tissue.



FASTER HEALING

StrataMGT promotes a moist healing environment leading to faster healing, and reduces the acute inflammatory response.



SYMPTOMATIC RELIEF

StrataMGT relieves symptoms such as itchiness, tenderness, dryness, burning sensation, painful intercourse (dyspareunia), painful urination (dysuria) and rectal or defecating pain.



RESTORES BARRIER FUNCTION

StrataMGT temporarily restores the barrier function of the mucosa.



HYDRATION AND GAS PERMEABILITY

StrataMGT is semi-permeable, which allows the mucosa to breathe and remain hydrated.



NON-REACTIVE

StrataMGT is inert, has no measurable pH, contains no hormones, steroids, alcohol, parabens or fragrances, and is suitable for long-term use.



NON-ABSORBABLE

When used inside the vagina or rectum, StrataMGT will not be absorbed systemically.

Directions for use



1. Wash your hands before applying StrataMGT.



 When used vaginally, apply
 to 5 drops of StrataMGT on the tip of your finger.*



3. Distribute StrataMGT evenly internally and externally as required.

Additional directions

- For best results StrataMGT should be maintained in continuous contact with the affected area (24 hours a day / 7 days a week).
- StrataMGT should be applied as needed or as advised by your physician.
- StrataMGT may be re-applied more often to ensure constant contact with mucosa or skin, or to reduce symptoms.
- StrataMGT should be applied after each urination, bowel movement or wash.
- StrataMGT does not need to be rubbed in or massaged, as it does not penetrate through mucosa or skin and will not enhance its effect.
- It is recommended to use panty liners during the first weeks of StrataMGT use, in case of product leaking.
- StrataMGT may stain clothing or sheets. If staining occurs, dry cleaning should be able to remove it without damaging the fabric.
- StrataMGT should not be applied over topical medications unless advised by the physician.

How much StrataMGT® do I need?



StrataMGT gel is an advanced formulation that requires **substantially less gel per application** than typical creams or gels.

StrataMGT 1 oz (28.4 g) contains enough gel for approximately 4 weeks of use 2-3 times per day.

Patients with different symptoms or diagnoses may require more-or-less product or numbers of applications per day.

Generally, patients will understand how much and how often to apply for their relief within a few days.

Duration of the treatment

StrataMGT should be applied for a minimum of 30 days.

StrataMGT is **indicated for short- and long-term use** to maintain the health of the mucosa, and should be continued as required.

StrataMGT® and other products

Moisturizers, lotions, etc. are not required. StrataMGT can be re-applied more often to avoid dry and tight skin feeling, as StrataMGT prevents the water evaporation through the damaged mucosa that may cause this feeling.



StrataMGT may reduce the effectiveness of contraception

such as spermicides, condoms or vaginal devices. Please use other precautions during this time.

StrataMGT **may increase lubrication** and therefore should not be used in conjunction with condoms for the prevention of sexually transmitted diseases (STDs).

^{*}Some patients prefer to use a common vaginal applicator for intravaginal application.



StrataMGT® - a non-hormonal and non-steroidal gel for vaginal atrophy and lichenoid dermatoses

- StrataMGT relieves symptoms such as itchiness, tenderness, dryness, burning sensation, painful intercourse (dyspareunia), painful urination (dysuria), rectal or defecating pain.
- StrataMGT improves erythema, mucosal tissue thinning, erosions, fissures, ulcerations, scarring/adhesions and swelling.
- StrataMGT is inert, has no measurable pH, contains no hormones, steroids, alcohol, parabens or fragrances, and is suitable for long-term use.
- StrataMGT is gas-permeable, allowing the vaginal mucosa to breathe and remain hydrated.
- · StrataMGT promotes faster healing.
- StrataMGT is easy to apply at home.











For any questions about StrataMGT or Stratpharma products, please contact us at our San Diego Headquarters:

619-930-5788 or customerservice@us.stratpharma.com Stratpharma Inc, 7676 Hazard Center Drive, Suite 880, San Diego, CA 92108 USA

us.stratamgt.com

Caution: For external use only. Consult instructions for use. Keep out of reach of children. For further instructions, please refer to the Patient Information Leaflet (PIL).

Ingredients: Polydimethylsiloxanes, siloxanes, alkylmethyl silicones

References: 1. Wysocki, 5. et al. (2014). Clinical medicine insights. Reproductive health, 8., pp. 23-30.

2. Bleibel, B., Nguyen, H. (2022). Vaginal Atrophy. In StatPearls. StatPearls Publishing. 3. Angelou, K. et al. (2020). Cureus, 12(4), e7586. 4. Palacios, S. et al. (2018). Climacteric: the Jot file International Menopause Society, 21(3), pp. 286-291. 5. Benini, V. et al. (2022). Medicina, 58(6), pp. 770. 6. Arroyo, (2017). International Jot Momers Health, 9, pp. 591-595. A chinkan, S. (2018). Haydrapsa Numune Med./, 58(2), pp.112-115. 8. Corazza, M. et al. (2021). Biomedicines, 9(8), pp. 950. 9. Data on file, 2022 (Orange Coast Womens Medical Group, USA). Stratpharma AG. 10. Data on file, 2022 (Orange Coast Womens Medical Group, USA). Stratpharma AG.

Manufactured by: Stratpharma AG, Aeschenvorstadt 57, CH-4051 Basel, Switzerland San Diego Headquarters: 619-930-5788, customerservice@us.stratpharma.com, Stratpharma Inc, 7676 Hazard Center Drive, Suite 880, San Diego, CA 92108 USA

